INTERNATIONAL JOURNAL OF QUANTUM CHEMISTRY, VOL. VIII, 999–1002 (1974)

Author Index, Volume VIII

Ali, M. K. and Meath, W. J.: Floating one-center perturbation treatments for H_2^+-like molecules based on screened hydrogen atom or molecular puff unperturbed problems, 119

Almlöf, A.: *Ab initio* calculations on porphrin, 915

Amos, A. T.: See Laughlin, C.

Aslangul, C., Constanciel, R., Daudel, R., Esnault, L. and Ludena, E. V.: The Loge theory as a starting point for variational calculations. I. General formalism, 499

Bandrauk, A. D.: Integral transform functions and separable potentials, 209

Basilevsky, M. V. and Berenfeld, M. M.: Interaction of two H_2 molecules; comparative study of various perturbation procedures, 467

Bendazzoli, G. I. and Palmieri, P.: Spin-orbit interaction in polyatomic molecules: *ab initio* computations with Gaussian orbitals, 941

Benston, M. L.: See Chong, D. P.

Berenfeld, M. M.: See Basilevsky, M. V.

Bhattacharya, A. K.: See Gupta, A.

Boyd, R. J. and Katriel, J.: The Coulomb hole in the 2^3S state of the helium isoelectronic sequence, 255

Boyle, L. L. and Schäffer, C. E.: Alternative equivalent icosahedral irreducible tensors, 153

Caballol, R., Gallifa, R., Riera, J. M. and Carbo, R.: Generalized open-shell SCF theory, 373

Cantu, A. A.: See Klein, D. J.

Carbo, R.: Eigenspace manipulation in SCF theory: general formalism, 423

Carbo, R.: See Caballol, R.

Chong, D. P., Scott, W. R., Yue, C. P., Wang, P. S. C., Benston, M. L. and Palke, W. E.: Application of the distinguishable electron method, 137

Čížek, J., Paldus, J. and Hubač, I.: Correlation effects in low-lying excited states of the PPP models of alternant hydrocarbons. I. Qualitative rules for the effect of limited configuration interaction, 951

Claverie, P.: See Daudey, J. P.

Coffey, P.: Potential energy integrals in semiempirical MO methods, 263

Coffey, P.: Virtual orbital transformation prior to configuration interaction with localized orbitals, 777

Constanciel, R.: See Aslangul, C.

Corsaro, F. A., Solenberger, F. R. and Hameka, H. F. A variational pertubation treatment of the hydrogen molecular ion using the Green function method on the united atom, 319

Damommio, A. and Synek, M.: Accurate analytical self-consistent field wave functions for Tm^{2+} and Tm^{3+}, 73

Davidson, E. R.: Matrix elements for spin-adapted configurations, 83

Davidson, E. R.: See Elbert, S.T.

Davidson, E. R.: See Jackels, C. F.

Davidson, E. R.: See Langhoff, S. R.

Daudel, R.: See Aslangul, C.

Daudey, J. P.: Perturbative *ab initio* calculations of intermolecular energies. III. The water dimer, 29

Daudey, J. P., Claverie, P. and Malrieu, J. P.: Perturbative *ab initio* calculations of intermolecular energies. I. Method, 1

Daudey, J. P., Malrieu, J. P. and Rojas, O.: Perturbative *ab initio* calculations of intermolecular energies. II. The He $\cdots$ He problem, 17

Delgado-Barrio, G.: See Smeyers, Y. G.

Delhalle, J.: On the need of precision in the calculation of the LCAO density matrix of polymers, 201

Dement'ev, A. I., Stepanov, N. F. and Yarovoi, S. S.: Convergence problems in the solution of SCF equations, 107

Domany, E.: See Katriel, J.

Elbert, S. T. and Davidson, E. R.: *Ab initio* calculations on urea, 857

Emanuel, R. V.: An investigation into the use of expansion methods in the calculation of chemical shifts and diamagnetic susceptibilities, 745
Esnault, L.: See Aslangul, C.

Fowler, D. R.: See Pohl, H. A.

Gallifa, R.: See Caballol, R.
Gallup, G. A.: Symmetries of n-electron wave functions under various spatial groups, 267
Garton, D.: Gradient of the general energy expression, 619
Grosswendt, B.: See Witschel, W.
Guerillot, C. R. and Lissillour, R.: Electronic correlation studies: II. Self-correlated field method. Application to ground state and first two ^1P, ^3P excited states of two-electron atomic systems, 825
Gupta, A., Bhattacharya, A. K. and Mukherjee, P. K.: Coupled Hartree–Fock calculation of static dipole polarizabilities and shielding factors of open-shell systems, 97

Hackmeyer, M.: A general *ab initio* molecular multi-configuration self-consistent field algorithm, 783
Hameka, H. F.: See Corsaro, F. A.
Hendeković, J.: Method of complex molecular orbitals, 799
Hameka, H. F.: See Svendsen, E. N.
Hubač, I.: See Čížek, J.

I'haya, Y. J.: See Narita, S.

Jackels, C. F. and Davidson, E. R.: Equivalence-restricted open-shell SCF theory, 707
Jansen, L.: See Lombardi, E.
Jucys, A. P. and Lazauskas, V. M.: Extended Hartree–Fock self-consistent field with some results for berylliumlike atomic systems, 839

Kagawa, T. and Murai, T.: Electron correlation study for two-electron atoms by simple correlated wave function, 755
Katriel, J., Domany, E.: A study of the symmetry dilemma: second-order transitions, 559
Katriel, J.: See Boyd, R. J.
Kiss, A. I. and Martin, A.: Molecules-in-molecules calculations with fixed resonance integrals, 427
Klein, D. J.: Comments on kets not strongly symmetry adapted, 235
Klein, D. J. and Cantu, A. A.: Symmetry adaptation and the relation between the spin-free and space-spin formulations of electronic structure, 223
König, E. and Kremer, S.: Irreducible tensor operator methods in intermediate-field coupling, 347
Kremer, S.: See König, E.
Krusius, P.: See von Boehm, J.

Langhoff, S. R. and Davidson, E. R.: Configuration interaction calculations on the nitrogen molecule, 61
Laughlin, C., Woodward, M. R. and Ammos, A. T.: Brillouin-Wigner perturbation theory and the generalized eigenvalue equation, 491
Lazauskas, V. M.: See Jucys, A. P.
Ledovskaya, E. M. and Petrashen, M. I.: Symmetry in the Heitler-London method. I. The connection of the molecular point group with its electron permutation group in the Heitler-London method, 163
Ledovskaya, E. M. and Petrashen, M. I.: Symmetry in the Heitler-London method. II. The determination of the allowed molecular multiplets, 171
Leibovici, C.: Structure de l'ion fluoronium FH_3^+ et chemin de la protonation du fluorure d'Hydrogene, 193
Letoquart, D.: Simplification of Kronecker products for the representations of point groups having only one n-fold axis with $n \geqq 3$, 627
Ley, B. E., Thorpe, T. and Rothstein, S. M.: Error analysis of variationally optimized upper- and lower-bound wave functions for H_4^+, 971
Lim, T. K.: Waller-Hartree spin-free method, 523
Lim, T. K.: SCF equations for pure spin states with many-particle interactions, 539
Linderberg, J. and Prato, D.: Dynamic polarizability of helium: A random phase approximation calculation, 901
Linderberg, J. and Seamans, L.: Matrix elements in all valence electron models, 925
Lissillour, R.: See Guerillot, C. R.
Lombardi, E., Pirola, L., Tarantini, G.,

Jansen, L. and Ritter, R.: Validity of the three-center, four-electron model for stability of rare gas halides on the basis of exchange perturbation theory, 335
Loubriel, G. M. and Selsby, R. G.: A technique for orbital exponent optimization in *ab initio* HF–SCF–LCAO–MO calculations, 547
Ludena, E. V.: See Aslangul, C.

Malrieu, J. P.: See Daudey, J. P.
Martin, A.: See Kiss, A. I.
Matcha, R. L. and Milleur, M. B.: Unified treatment of two-electron integrals. II. One-center, electron repulsion, spin-spin and spin-orbit intergrals containing r_{ij}, 635
Mathur, S. C. and Singh, J.: More on molecular electronic structure of metal phthalocyanines, 79
Mayer, I.: Spin projected EHF method: calculations for a four-electron model system, 363
Mayer, I.: Spin-projected extended Hartree–Fock equations. II. Odd-electron systems, 893
Meath, W. J.: See Ali, M. K.
Mestechkin, M. M. and Whyman, G. E.: Matrix formulation of the generalized Hartree–Fock methods, 45
Milleur, M. B.: See Matcha, R. L.
Moccia, R.: Static and dynamic first- and second-order properties by variational wave functions, 293
Mukherjee, D.: A comparative calculation on excited state energies of some conjugated hydrocarbons, 247
Mukherjee, P. K.: See Gupta, A.
Mulliken, R. S.: Molecular orbitals of nitrogen at small internuclear distances, 817
Murai, T.: See Kagawa, T.

Nanda, D. N. and Narasimhan, P. T.: Approximate valence-shell electron unrestricted Hartree-Fock calculations with orthogonalized atomic orbitals: proton hyperfine splittings in benzyl and related radicals, 451
Narasimhan, P. T.: See Nanda, D. N.
Narita, S. and I'haya, Y. J.: Dual-basis orthonormality-constrained variation method, 765

Paldus, J.: See Čížek, J.
Palke, W. E.: See Chong, D. P.
Palmieri, P.: See Bendazzoli, G. I.
Petrashen, M. I.: See Ledovskaya, E. M.
Pirola, L.: See Lombardi, E.
Pohl, H. A. and Fowler, D. R.: An improved Hellmann-type pseudopotential for atoms and molecules, 435
Prato, D.: See Linderberg, J.
Pulfer, J. D. and Whitehead, M. A.: Hammerstein integral equivalent of Ricatti's equation, 677
Pulfer, J. D. and Whitehead, M. A.: The Hammerstein integral equation: a general technique for constructing a rapidly convergent Padé-type approximation of the logarithmic derivative, 693

Randić, M.: Hybridization by the maximum overlap method, 643
Riera, J. M.: See Caballol, R.
Ritter, R.: See Lombardi, E.
Rojas, O.: See Daudey, J. P.
Rothstein, S. M.: See Ley, B. E.

Sasaki, F.: Matrix elements in configuration interaction calculations, 605
Schäffer, C. E.: See Boyle, L. L.
Scott, W. R.: See Chong, D. P.
Seamans, L.: See Linderberg, J.
Selsby, R. G.: See Loubriel, G. M.
Singh, J.: See Mathur, S. C.
Smeyers, Y. G. and Delgado-Barrio, G.: Half-projected and projected Hartree-Fock calculations for singlet ground states. II. Lythium hyride, 733
Solenberger, F. R.: See Corsaro, F. A.
Stenger, W.: Intermediate problems for eigenvalues, 623
Stepanov, N. F.: See Dement'ev, A. I.
Sundaram, K.: Nonbonded interactions in membrane active cyclic biopolymers. I. Valinomycin-potassium ion complex, 565
Svendsen, E. N. and Hameka, H. F.: Calculation of the magneto-optical activity of the hydrogen molecule, 789
Synek, M.: See Damommio, A.

Tanaka, K.: Basis dependence of wave functions of the ground state and low-lying excited states of planar formaldehyde, 981
Tarantini, G.: See Lombardi, E.

Thorpe, T.: See Ley, B. E.

Wang, P. S. C.: See Chong, D. P.
Whitehead, M. A.: See Pulfer, J. D.
Witschel, W. and Grosswendt, B.: Oscillator and hydrogenic matrix elements by operator algebra, 585
Whyman, G. E.: See Mestechkin, M. M.
Woodward, M. R.: See Laughlin, C.
Wormer, P. E. S. and Van Der Avoird, A.: Transformation properties of many-electron wave functions with special attention to the relation between pair-correlated DODS and configuration interaction, 715

Van Der Avord, A.: See Wormer, P. E. S.
von Boehm, J. and Krusius, P.: The self-consistent symmetrized OPW method with an application to crystalline selenium, 395
van Duijnen, P. Th.: Molecular integrals over generalized Gaussian basis sets, 179

Yarovoi, S. S.: See Dement'ev, A. I.
Yue, C. P.: See Chong, D. P.

Zeroka, D.: Variation of the polarizability of the hydrogen molecule ion and the hydrogen molecule with internuclear separation, 91